

# Package ‘ensemblepp’

October 13, 2022

**Title** Ensemble Postprocessing Data Sets

**Version** 1.0-0

**Date** 2019-05-03

**Author** Jakob Messner [aut, cre]

**Maintainer** Jakob Messner <jakob.messner@posteo.net>

**Depends** R (>= 2.10.0)

**Suggests** ensembleBMA, crch, gamlss, ensembleMOS, SpecsVerification,  
scoringRules, glmx, ordinal, pROC, mvtnorm

**Description** Data sets for the chapter ``Ensemble Postprocessing with R'' of the book Stephane Vanitsem, Daniel S. Wilks, and Jakob W. Messner (2018) ``Statistical Postprocessing of Ensemble Forecasts'', Elsevier, 362pp. These data sets contain temperature and precipitation ensemble weather forecasts and corresponding observations at Innsbruck/Austria. Additionally, a demo with the full code of the book chapter is provided.

**License** GPL-2 | GPL-3

**NeedsCompilation** no

**Repository** CRAN

**Date/Publication** 2019-05-08 07:50:10 UTC

## R topics documented:

rain . . . . .	2
temp . . . . .	3

**Index**

**5**

rain

*Precipitation Observations and Forecasts for Innsbruck*

## Description

Accumulated 18-30 hour precipitation ensemble forecasts and corresponding observations at Innsbruck. The dataset includes GEFS reforecasts (Hamill et al. 2013) and observations from SYNOP station Innsbruck Airport (11120) from 2000-01-02 to 2016-01-01.

## Usage

```
data("temp")
```

## Format

A data frame with 2749 rows. The first column (`rain`) are 12-hour accumulated precipitation observations. Columns 2-12 (`rainfc`) are 18-30 hour accumulated precipitation forecasts from the individual ensemble members.

## Source

Observations: <http://www.ogimet.com/synops.phtml.en>

Reforecasts: <http://www.esrl.noaa.gov/psd/forecasts/reforecast2/>

## References

Hamill TM, Bates GT, Whitaker JS, Murray DR, Fiorino M, Galarneau Jr TJ, Zhu Y, Lapenta W (2013). NOAA's Second-Generation Global Medium-Range Ensemble Reforecast Data Set. *Bulletin of the American Meteorological Society*, 94(10), 1553-1565.

Vannitsem S, Wilks DS, Messner JW (2017). Statistical Postprocessing of Ensemble Forecasts, Elsevier, to appear.

## Examples

```
## Diagnostic plots similar to Figure 8 in Vannitsem et al. ##

## load and prepare data
data("rain")

rain <- sqrt(rain)

rain$ensmean <- apply(rain[,2:12], 1, mean)
rain$enssd <- apply(rain[,2:12], 1, sd)

## Scatterplot of precipitation by ensemble mean
plot(rain$ensmean, rain, col = gray(0.2, alpha = 0.4),
     main = "Scatterplot")
abline(0, 1, lty = 2)
```

```
## Verification rank histogram
rank <- apply(rain[,1:12], 1, rank)[1,]
hist(rank, breaks = 0:12 + 0.5, main = "Verification Rank Histogram")

## Spread skill relationship
sdcat <- cut(rain$enssd, quantile(rain$enssd, seq(0, 1, 0.2)))
boxplot(abs(rain$ensmean)~sdcat, rain, ylab = "absolute error",
xlab = "ensemble standard deviation", main = "Spread-Skill")

## Histogram
hist(rain$rain, xlab = "square root of precipitation", main = "Histogram")
```

---

temp

*Minimum Temperature Observations and Forecasts for Innsbruck*

---

## Description

18-30 hour minimum temperature ensemble forecasts and corresponding observations at Innsbruck. The dataset includes GEFS reforecasts (Hamill et al. 2013) and observations from the SYNOP station Innsbruck Airport (11120) from 2000-01-02 to 2016-01-01.

## Usage

```
data("temp")
```

## Format

A data frame with 2749 rows. The first column (`temp`) are 12-hour minimum temperature observations. Columns 2-12 (`tempfc`) are 18-30 hour minimum temperature forecasts from the individual ensemble members.

## Source

Observations: <http://www.ogimet.com/synops.phtml.en>

Reforecasts: <http://www.esrl.noaa.gov/psd/forecasts/reforecast2/>

## References

Hamill TM, Bates GT, Whitaker JS, Murray DR, Fiorino M, Galarneau Jr TJ, Zhu Y, Lapenta W (2013). NOAA's Second-Generation Global Medium-Range Ensemble Reforecast Data Set. *Bulletin of the American Meteorological Society*, 94(10), 1553-1565.

Vannitsem S, Wilks DS, Messner JW (2017). Statistical Postprocessing of Ensemble Forecasts, *Elsevier*, to appear.

## Examples

```
## Diagnostic plots similar to Figure 1 and 3 in Vannitsem et al. ##  
  
## load and prepare data  
data("temp")  
  
temp$ensmean <- apply(temp[,2:12], 1, mean)  
temp$enssd <- apply(temp[,2:12], 1, sd)  
  
## Scatterplot of minimum temperature observation by ensemble mean  
plot(temp~ensmean, temp, main = "Scatterplot")  
abline(0, 1, lty = 2)  
  
## Verification rank histogram  
rank <- apply(temp[,1:12], 1, rank)[1,]  
hist(rank, breaks = 0:12 + 0.5, main = "Verification Rank Histogram")  
  
## Spread skill relationship  
sdcat <- cut(temp$enssd, breaks = quantile(temp$enssd, seq(0, 1, 0.2)))  
boxplot(abs(temp-ensmean)~sdcat, temp, ylab = "absolute error",  
xlab = "ensemble standard deviation", main = "Spread-Skill")  
  
## Histogram  
hist(temp$temp, xlab = "minimum temperature", main = "Histogram")
```

# Index

\* **datasets**

    rain, [2](#)

    temp, [3](#)

    rain, [2](#)

    temp, [3](#)